

MI Preprint Series

Kyushu University
The Global COE Program
Math-for-Industry Education & Research Hub

On very accurate enclosure of the
optimal constant in the a priori
error estimates for H_0^2 -projection

T. Kinoshita & M. T. Nakao

MI 2009-25

(Received August 5, 2009)

Faculty of Mathematics
Kyushu University
Fukuoka, JAPAN

On very accurate enclosure of the optimal constant in the a priori error estimates for H_0^2 -projection

Takehiko Kinoshita*, Mitsuhiro T. Nakao

Faculty of Mathematics, Kyushu University, Hakozaki, Higashiku, Fukuoka 812-8581, Japan

Abstract

We present constructive a priori error estimates for H_0^2 -projection into a space of polynomials on a one-dimensional interval. Here, “constructive” indicates that we can obtain the error bounds in which all constants are explicitly given or are represented in a numerically computable form. Using the properties of Legendre polynomials, we consider a method by which to determine these constants to be as small as possible. Using the proposed technique, the optimal constant could be enclosed in a very narrow interval with results verification. Furthermore, constructive error estimates for finite element H_0^2 -projection in one dimension are presented. This type of estimates will play an important role in the numerical verification of solutions for nonlinear fourth-order elliptic problems as well as in the guaranteed a posteriori error analysis for the finite element method or the spectral method.

Key words: Constructive a priori error estimates, Legendre polynomials, Fourth-order elliptic problem

PACS: 02.60.Ed, 02.60.Lj, 02.70.Dh, 02.70.Hm

2000 MSC: 65L10, 65L60, 65L70, 65N15, 65N30, 65N35, 65G99, 35J40

*Corresponding author

Email addresses: t-kinoshita@math.kyushu-u.ac.jp (Takehiko Kinoshita), mtanakao@math.kyushu-u.ac.jp (Mitsuhiro T. Nakao)

1. Introduction

In the present paper, we consider the smallest constant C in a priori error estimates of the form

$$\|u - P^2 u\|_{H_0^2(\Lambda)} \leq C |u|_{H^4(\Lambda)}, \quad \forall u \in H_0^2(\Lambda) \cap H^4(\Lambda), \quad (1)$$

where P^2 is an H_0^2 -projection on a one-dimensional interval Λ , and $\|\cdot\|_{H_0^2(\Lambda)}$ and $|\cdot|_{H^4(\Lambda)}$ are the norm in H_0^2 and the seminorm in H^4 , respectively. The purpose of the present study is to find the upper and lower bounds of optimal constants in the above estimates. These constants not only play an important role in theoretically verifying the solutions of differential equations (e.g. [8, 2, 6]), but also contribute to highly reliable computing in numerical simulation using the finite element method or the spectral method. In general, C should be made as small as possible.

In the case of the H_0^1 -projection, for approximation spaces with linear and quadratic polynomials, the optimal constants can be theoretically determined as $\frac{1}{\pi}$ and $\frac{1}{2\pi}$, respectively (see [7, 5]). Such a constant can also be computed for higher-order polynomials (see [4]), although it is not optimal.

For the H_0^2 -projection, Schultz obtained constructive a priori error estimates based on piecewise cubic interpolation (see [7]), which is not optimal.

In the present paper, we propose a method that is an extension and improvement of the technique presented in [4] to obtain a constant very close to the optimal constant with guaranteed accuracy. Note that the proposed technique improves Schultz's result and can also be applied to obtain the optimal constants in the case of higher-order polynomials. Furthermore, using the present results, it will be possible to realize more efficient computations in the numerical verification of solutions related to fourth-order elliptic problems, such as those described in [2, 6, 9].

2. Legendre polynomials

Let $\Lambda = (a, b)$, ($a < b \in \mathbb{R}$) be a one-dimensional interval. The Legendre polynomials on Λ are defined as a complete orthogonal system in $L^2(\Lambda)$, for an arbitrary non-negative integer n ,

$$P_n(x) := \frac{(-1)^n}{n! |\Lambda|^n} \left(\frac{d}{dx} \right)^n (b-x)^n (x-a)^n, \quad (2)$$

where $|\Lambda| := b - a$. Furthermore, P_n has the following properties ([1]):

$$\frac{d}{dx} ((b-x)(x-a)P'_n(x)) + n(n+1)P_n(x) = 0, \quad \forall n \geq 0, \quad (3)$$

$$(P_m, P_n)_{L^2(\Lambda)} = \frac{|\Lambda|}{2n+1} \delta_{m,n}, \quad \forall m, n \geq 0, \quad (4)$$

$$(2n+1)P_n = \frac{|\Lambda|}{2} (P'_{n+1} - P'_{n-1}), \quad \forall n \geq 1, \quad (5)$$

$$P_n(a) = (-1)^n, \quad P_n(b) = 1, \quad \forall n \geq 0, \quad (6)$$

where $(P_m, P_n)_{L^2(\Lambda)}$ denotes the L^2 inner product on Λ and $\delta_{m,n}$ denotes Kronecker's δ .

Lemma 2.1. *For any $u \in H^1(\Lambda)$ and integer $n \geq 1$, we have*

$$(u, P_n)_{L^2(\Lambda)} = \frac{|\Lambda|}{2(2n+1)} \left((u', P_{n-1})_{L^2(\Lambda)} - (u', P_{n+1})_{L^2(\Lambda)} \right). \quad (7)$$

Proof: From (5) and (6)

$$\begin{aligned} (u, P_n)_{L^2(\Lambda)} &= \left(u, \frac{|\Lambda|}{2(2n+1)} (P'_{n+1} - P'_{n-1}) \right)_{L^2(\Lambda)} \\ &= \frac{|\Lambda|}{2(2n+1)} u(b) (P_{n+1}(b) - P_{n-1}(b)) - \frac{|\Lambda|}{2(2n+1)} u(a) (P_{n+1}(a) - P_{n-1}(a)) \\ &\quad - \frac{|\Lambda|}{2(2n+1)} \left((u', P_{n+1} - P_{n-1})_{L^2(\Lambda)} \right) \\ &= \frac{|\Lambda|}{2(2n+1)} \left((u', P_{n-1})_{L^2(\Lambda)} - (u', P_{n+1})_{L^2(\Lambda)} \right), \end{aligned}$$

which implies (7). \square

3. Error estimates for H_0^2 -projection on a one-dimensional interval

Let $H_0^2(\Lambda)$ be a function space on Λ defined as

$$H_0^2(\Lambda) \equiv \{u \in H^2(\Lambda) ; u(a) = u(b) = u'(a) = u'(b) = 0\}$$

with associated inner product

$$(u, v)_{H_0^2(\Lambda)} := (u'', v'')_{L^2(\Lambda)}.$$

First, we define the following set of functions

Definition 3.1. For any integer $n \geq 4$, an n -th order polynomial ϕ_n on Λ is defined as

$$\phi_n(x) = \frac{(-1)^n \sqrt{2n-3}}{(n-2)! |\Lambda|^{n-3/2}} \left(\frac{d}{dx} \right)^{n-4} (b-x)^{n-2} (x-a)^{n-2}. \quad (8)$$

Then, we have

Theorem 3.2. The set of functions $\{\phi_n\}_{n \geq 4} \subset H_0^2(\Lambda)$ is a complete orthonormal system in $H_0^2(\Lambda)$.

Proof: First, we show the orthogonality. From (8) we have, for arbitrary $n \geq 2$,

$$\begin{aligned} \phi_{n+2}''(x) &= \sqrt{\frac{2n+1}{|\Lambda|}} \frac{(-1)^n}{n! |\Lambda|^n} \left(\frac{d}{dx} \right)^n (b-x)^n (x-a)^n \\ &= \sqrt{\frac{2n+1}{|\Lambda|}} P_n(x). \end{aligned}$$

Hence, for any $m, n \geq 2$, by using the property given in (4), it holds that

$$\begin{aligned} (\phi_{m+2}, \phi_{n+2})_{H_0^2(\Lambda)} &= (\phi_{m+2}'', \phi_{n+2}'')_{L^2(\Lambda)} \\ &= \left(\sqrt{\frac{2m+1}{|\Lambda|}} P_m, \sqrt{\frac{2n+1}{|\Lambda|}} P_n \right)_{L^2(\Lambda)} \\ &= \frac{\sqrt{(2m+1)(2n+1)}}{|\Lambda|} (P_m, P_n)_{L^2(\Lambda)} \\ &= \delta_{m,n}, \end{aligned}$$

which implies that $\{\phi_n\}_{n \geq 4}$ is an orthonormal system in $H_0^2(\Lambda)$.

Next, we prove the completeness. For an element $u \in H_0^2(\Lambda)$, suppose that $(u, \phi_{n+2})_{H_0^2(\Lambda)} = 0$ for all $n \geq 2$. Then, we have

$$(u, \phi_{n+2})_{H_0^2(\Lambda)} = \sqrt{\frac{2n+1}{|\Lambda|}} (u'', P_n)_{L^2(\Lambda)}, \quad \forall n \geq 2.$$

Namely,

$$(u'', P_n)_{L^2(\Lambda)} = 0, \quad \forall n \geq 2.$$

Moreover, by $u \in H_0^2(\Lambda)$ we have the following equalities:

$$\begin{aligned}(u'', P_0)_{L^2(\Lambda)} &= u'(b) - u'(a) = 0, \\ (u'', P_1)_{L^2(\Lambda)} &= -\frac{2}{|\Lambda|} (u', P_0)_{L^2(\Lambda)} = -\frac{2}{|\Lambda|} (u(b) - u(a)) = 0.\end{aligned}$$

Since $\{P_n\}_{n \geq 0}$ is a complete orthogonal system in $L^2(\Lambda)$, it holds that $u'' = 0$ in $L^2(\Lambda)$. Thus, we have $u = 0$ in $H_0^2(\Lambda)$, which proves the completeness of $\{\phi_n\}_{n \geq 4}$. \square

Definition 3.3 (H_0^2 -projection). For an integer $N \geq 4$, we define the finite-dimensional subspace S_N of $H_0^2(\Lambda)$ by $S_N \equiv \text{span}_{4 \leq n \leq N} \phi_n$. Then, we define the H_0^2 -projection P_N^2 from $H_0^2(\Lambda)$ into S_N by

$$(u - P_N^2 u, v_N)_{H_0^2(\Lambda)} = 0, \quad \forall v_N \in S_N. \quad (9)$$

We also set $S_3 = \{0\}$ and $P_3^2 \equiv 0$.

Now, we have the following basic constructive error estimates for the H_0^2 -projection of a function u with H^4 -regularity.

Theorem 3.4. For an arbitrary integer $N \geq 3$, there exists a constant $\tilde{C}(|\Lambda|, N) > 0$ such that

$$\|u - P_N^2 u\|_{H_0^2(\Lambda)} \leq \tilde{C}(|\Lambda|, N) |u|_{H^4(\Lambda)}, \quad \forall u \in H_0^2(\Lambda) \cap H^4(\Lambda), \quad (10)$$

where

$$\tilde{C}(|\Lambda|, N) = \begin{cases} \sqrt{3} \left(\frac{|\Lambda|}{2}\right)^2 \frac{1}{\sqrt{2N-5}(2N-3)\sqrt{2N-1}}, & \text{if } N = 3, \\ \sqrt{3} \left(\frac{|\Lambda|}{2}\right)^2 \frac{\sqrt{10N-3}}{(2N-3)\sqrt{(2N-1)(2N+1)(2N+3)}}, & \text{if } 4 \leq N \leq 38, \\ 3\sqrt{2} \left(\frac{|\Lambda|}{2}\right)^2 \frac{1}{\sqrt{(2N-1)(2N+1)(2N+5)(2N+7)}}, & \text{if } 39 \leq N. \end{cases}$$

Here, the H^4 seminorm is defined as $|u|_{H^4(\Lambda)} \equiv \|u''''\|_{L^2(\Lambda)}$.

Proof: From Theorem 3.2, any $u \in H_0^2(\Lambda) \cap H^4(\Lambda)$ can be expanded by $\{\phi_n\}$. That is,

$$u = \sum_{n=4}^{\infty} a_n \phi_n \quad (11)$$

$$\text{with } a_n = (u, \phi_n)_{H_0^2(\Lambda)}. \quad (12)$$

As a result of the orthogonality of $\{\phi_n\}$ in $H_0^2(\Lambda)$, the H_0^2 -projection coincides with the truncation up to N . Hence, we have

$$P_N^2 u = \sum_{n=4}^N a_n \phi_n.$$

Therefore, the Parseval equality implies the following:

$$\begin{aligned} \|u - P_N^2 u\|_{H_0^2(\Lambda)}^2 &= \left\| \sum_{n=N+1}^{\infty} a_n \phi_n \right\|_{H_0^2(\Lambda)}^2 \\ &= \sum_{n=N+1}^{\infty} a_n^2. \end{aligned} \quad (13)$$

On the other hand, since $\{P_n\}$ is a complete orthogonal system in $L^2(\Lambda)$, $u'''' \in L^2(\Lambda)$ can also be expanded as

$$u'''' = \sum_{n=0}^{\infty} b_n \frac{P_n}{\|P_n\|_{L^2(\Lambda)}} \quad (14)$$

$$\text{with } b_n = \left(u'''' , \frac{P_n}{\|P_n\|_{L^2(\Lambda)}} \right)_{L^2(\Lambda)}. \quad (15)$$

Taking into account that $P_n / \|P_n\|_{L^2(\Lambda)}$ is a complete orthonormal system in $L^2(\Lambda)$, by the Parseval equality, we have

$$|u|_{H^4(\Lambda)}^2 = \|u''''\|_{L^2(\Lambda)}^2 = \sum_{n=0}^{\infty} b_n^2. \quad (16)$$

Now, for any integer $n \geq 4$, observe that by using Lemma 2.1

$$\begin{aligned}
a_n &= (u, \phi_n)_{H_0^2(\Lambda)} \\
&= (u'', \phi_n'')_{L^2(\Lambda)} \\
&= \sqrt{2n-3} |\Lambda|^{-1/2} (u'', P_{n-2})_{L^2(\Lambda)} \\
&= \frac{|\Lambda|^{1/2}}{2\sqrt{2n-3}} (u''', P_{n-3} - P_{n-1})_{L^2(\Lambda)} \\
&= \frac{|\Lambda|^{1/2}}{2\sqrt{2n-3}} \frac{|\Lambda|}{2(2n-5)} (u'''' , P_{n-4} - P_{n-2})_{L^2(\Lambda)} - \frac{|\Lambda|^{1/2}}{2\sqrt{2n-3}} \frac{|\Lambda|}{2(2n-1)} (u'''' , P_{n-2} - P_n)_{L^2(\Lambda)} \\
&= \frac{|\Lambda|^{3/2}}{4} \frac{1}{(2n-5)\sqrt{2n-3}} (u'''' , P_{n-4})_{L^2(\Lambda)} - \frac{|\Lambda|^{3/2}}{4} \frac{2\sqrt{2n-3}}{(2n-5)(2n-1)} (u'''' , P_{n-2})_{L^2(\Lambda)} \\
&\quad + \frac{|\Lambda|^{3/2}}{4} \frac{1}{\sqrt{2n-3}(2n-1)} (u'''' , P_n)_{L^2(\Lambda)} \\
&=: \left(\frac{|\Lambda|}{2}\right)^2 (\alpha_{n-4} b_{n-4} - \beta_{n-2} b_{n-2} + \gamma_n b_n). \tag{17}
\end{aligned}$$

Here, α_n , β_n , and γ_n are defined, respectively, as follows:

$$\alpha_{n-4} = \frac{\|P_{n-4}\|_{L^2(\Lambda)}}{\sqrt{|\Lambda|(2n-5)\sqrt{2n-3}}} = \frac{1}{\sqrt{2n-7}(2n-5)\sqrt{2n-3}},$$

$$\text{namely, } \alpha_n = \frac{1}{\sqrt{2n+1}(2n+3)\sqrt{2n+5}}, \tag{18}$$

$$\beta_{n-2} = \frac{2\sqrt{2n-3} \|P_{n-2}\|_{L^2(\Lambda)}}{\sqrt{|\Lambda|(2n-5)(2n-1)}} = \frac{2}{(2n-5)(2n-1)},$$

$$\text{namely, } \beta_n = \frac{2}{(2n-1)(2n+3)}, \tag{19}$$

$$\gamma_n = \frac{\|P_n\|_{L^2(\Lambda)}}{\sqrt{|\Lambda|\sqrt{2n-3}(2n-1)}} = \frac{1}{\sqrt{2n-3}(2n-1)\sqrt{2n+1}}. \tag{20}$$

Note that α_n , β_n , and $\gamma_n \approx O(n^{-2})$ and are monotonically decreasing sequences in n . Then, we obtain the following estimates for each term of the

nal equality in (13)

$$\begin{aligned}
a_n^2 &= \left(\frac{|\Lambda|}{2}\right)^4 (\alpha_{n-4}b_{n-4} - \beta_{n-2}b_{n-2} + \gamma_n b_n)^2 \\
&= \left(\frac{|\Lambda|}{2}\right)^4 (\alpha_{n-4}^2 b_{n-4}^2 + \beta_{n-2}^2 b_{n-2}^2 + \gamma_n^2 b_n^2 \\
&\quad - 2\alpha_{n-4}b_{n-4}\beta_{n-2}b_{n-2} - 2\beta_{n-2}b_{n-2}\gamma_n b_n + 2\alpha_{n-4}b_{n-4}\gamma_n b_n) \\
&\leq 3 \left(\frac{|\Lambda|}{2}\right)^4 (\alpha_{n-4}^2 b_{n-4}^2 + \beta_{n-2}^2 b_{n-2}^2 + \gamma_n^2 b_n^2).
\end{aligned}$$

Therefore, from (13), we have the estimates

$$\begin{aligned}
\|u - P_N^2 u\|_{H_0^2(\Lambda)}^2 &= \sum_{n=N+1}^{\infty} a_n^2 \\
&\leq 3 \left(\frac{|\Lambda|}{2}\right)^4 \sum_{n=N+1}^{\infty} (\alpha_{n-4}^2 b_{n-4}^2 + \beta_{n-2}^2 b_{n-2}^2 + \gamma_n^2 b_n^2) \\
&= 3 \left(\frac{|\Lambda|}{2}\right)^4 \left(\alpha_{N-3}^2 b_{N-3}^2 + \alpha_{N-2}^2 b_{N-2}^2 + (\alpha_{N-1}^2 + \beta_{N-1}^2) b_{N-1}^2 \right. \\
&\quad \left. + (\alpha_N^2 + \beta_N^2) b_N^2 + \sum_{n=N+1}^{\infty} (\alpha_n^2 + \beta_n^2 + \gamma_n^2) b_n^2 \right) \\
&\leq 3 \left(\frac{|\Lambda|}{2}\right)^4 \max\{\alpha_{N-3}^2, \alpha_{N-1}^2 + \beta_{N-1}^2, \alpha_{N+1}^2 + \beta_{N+1}^2 + \gamma_{N+1}^2\} \sum_{n=N-3}^{\infty} b_n^2 \\
&\leq 3 \left(\frac{|\Lambda|}{2}\right)^4 \max\{\alpha_{N-3}^2, \alpha_{N-1}^2 + \beta_{N-1}^2, \alpha_{N+1}^2 + \beta_{N+1}^2 + \gamma_{N+1}^2\} |u|_{H^4(\Lambda)}^2 \\
&=: \tilde{C}(|\Lambda|, N)^2 |u|_{H^4(\Lambda)}^2.
\end{aligned}$$

Finally, estimating the terms operated on by $\max\{\dots\}$ in the above expression, we obtain for $N = 3$ then $\alpha_{N-1}^2 + \beta_{N-1}^2 \leq \alpha_{N-3}^2$ and $\alpha_{N+1}^2 + \beta_{N+1}^2 + \gamma_{N+1}^2 \leq \alpha_{N-3}^2$, which implies the following:

$$\tilde{C}(|\Lambda|, N) = \sqrt{3} \left(\frac{|\Lambda|}{2}\right)^2 \frac{1}{\sqrt{2N-5}(2N-3)\sqrt{2N-1}},$$

for $4 \leq N \leq 38$. Thus, $\alpha_{N-3}^2 \leq \alpha_{N-1}^2 + \beta_{N-1}^2$ and $\alpha_{N+1}^2 + \beta_{N+1}^2 + \gamma_{N+1}^2 \leq$

$\alpha_{N-1}^2 + \beta_{N-1}^2$, which implies the following:

$$\tilde{C}(|\Lambda|, N) = \sqrt{3} \left(\frac{|\Lambda|}{2} \right)^2 \frac{\sqrt{10N-3}}{(2N-3)\sqrt{(2N-1)(2N+1)(2N+3)}},$$

for $39 \leq N$ then $\alpha_{N-3}^2 \leq \alpha_{N+1}^2 + \beta_{N+1}^2 + \gamma_{N+1}^2$ and $\alpha_{N+1}^2 + \beta_{N+1}^2 + \gamma_{N+1}^2$, which implies

$$\tilde{C}(|\Lambda|, N) = 3\sqrt{2} \left(\frac{|\Lambda|}{2} \right)^2 \frac{1}{\sqrt{(2N-1)(2N+1)(2N+5)(2N+7)}}$$

Thus, we have the desired result. \square

Based on the estimates in Theorem 3.4, we can obtain a smaller constant by using a method similar to that described in [4].

Lemma 3.5. *Let $a_n, b_n \in \mathbb{R}$ be as given in the proof of Theorem 3.4. In addition, let $\alpha_n, \beta_n, \gamma_n$ be positive numbers defined by (18), (19), and (20), respectively. Then, for any integers $N \geq 3$ and $M \geq N + 5$, there exists a constant $\sigma_{N,M} > 0$ such that*

$$\sum_{n=N+1}^M a_n^2 \leq \left(\frac{|\Lambda|}{2} \right)^4 \sigma_{N,M} \sum_{n=N-3}^M b_n^2, \quad (21)$$

where $\sigma_{N,M} := \max\{c_1(N), c_2(N), c_3(N), d_1(M), d_2(M)\}$ and $c_i(N), d_i(M)$ are defined as follows:

$$\begin{aligned} c_1(N) &= \alpha_{N-3}^2 + \alpha_{N-3}\beta_{N-1} + \alpha_{N-3}\gamma_{N+1}, \\ c_2(N) &= \alpha_{N-3}\beta_{N-1} + \alpha_{N-1}^2 + \beta_{N-1}^2 + \alpha_{N-1}\beta_{N+1} + \beta_{N-1}\gamma_{N+1} + \alpha_{N-1}\gamma_{N+3}, \\ c_3(N) &= \alpha_{N-3}\gamma_{N+1} + \alpha_{N-1}\beta_{N+1} + \beta_{N-1}\gamma_{N+1} \\ &\quad + \alpha_{N+1}^2 + \beta_{N+1}^2 + \gamma_{N+1}^2 + \alpha_{N+1}\beta_{N+3} + \beta_{N+1}\gamma_{N+3} + \alpha_{N+1}\gamma_{N+5}, \\ d_1(M) &= \alpha_{M-7}\gamma_{M-3} + \alpha_{M-5}\beta_{M-3} + \beta_{M-5}\gamma_{M-3} + \beta_{M-3}^2 + \gamma_{M-3}^2 + \beta_{M-3}\gamma_{M-1}, \\ d_2(M) &= \alpha_{M-5}\gamma_{M-1} + \beta_{M-3}\gamma_{M-1} + \gamma_{M-1}^2. \end{aligned}$$

Proof: Setting $\vec{b} \equiv (b_{N-3}, b_{N-2}, \dots, b_M)^T \in \mathbb{R}^{M-N+4}$ and taking (17) into account, reveals that there exists a symmetric and positive definite matrix A satisfying

$$\begin{aligned} \sum_{n=N+1}^M a_n^2 &= \left(\frac{|\Lambda|}{2} \right)^4 \sum_{n=N+1}^M (\alpha_{n-4}b_{n-4} - \beta_{n-2}b_{n-2} + \gamma_n b_n)^2 \\ &= \left(\frac{|\Lambda|}{2} \right)^4 \vec{b}^T A \vec{b}. \end{aligned} \quad (22)$$

Theorem 3.6. For each integer $N \geq 3$, there exists a constant $C_0(|\Lambda|, N) > 0$ such that

$$\|u - P_N^2 u\|_{H_0^2(\Lambda)} \leq C_0(|\Lambda|, N) |u|_{H^4(\Lambda)}, \quad \forall u \in H_0^2(\Lambda) \cap H^4(\Lambda), \quad (23)$$

where

$$C_0(|\Lambda|, N) = \begin{cases} \left(\frac{|\Lambda|}{2}\right)^2 \sqrt{c_1(N)}, & \text{if } N = 3, \\ \left(\frac{|\Lambda|}{2}\right)^2 \sqrt{c_2(N)}, & \text{if } 4 \leq N \leq 19, \\ \left(\frac{|\Lambda|}{2}\right)^2 \sqrt{c_3(N)}, & \text{if } 20 \leq N, \end{cases}$$

where $c_i(N)$ are the constants given in Lemma 3.5.

Moreover, $c_i(N)$ are explicitly written as

$$\begin{aligned} c_1(N) &= \frac{1}{(2N-5)(2N-3)^2(2N-1)} + \frac{2}{\sqrt{2N-5}(2N-3)^2\sqrt{2N-1}(2N+1)} \\ &\quad + \frac{1}{\sqrt{2N-5}(2N-3)(2N-1)(2N+1)\sqrt{2N+3}}, \\ c_2(N) &= \frac{2}{\sqrt{2N-5}(2N-3)^2\sqrt{2N-1}(2N+1)} + \frac{4}{(2N-3)\sqrt{2N-1}(2N+1)\sqrt{2N+3}(2N+5)} \\ &\quad + \frac{1}{\sqrt{2N-1}(2N+1)(2N+3)(2N+5)\sqrt{2N+7}} + \frac{10N-3}{(2N-3)^2(2N-1)(2N+1)(2N+3)}, \\ c_3(N) &= \frac{1}{\sqrt{2N-5}(2N-3)(2N-1)(2N+1)\sqrt{2N+3}} + \frac{4}{(2N-3)\sqrt{2N-1}(2N+1)\sqrt{2N+3}(2N+5)} \\ &\quad + \frac{6}{(2N-1)(2N+1)(2N+5)(2N+7)} + \frac{4}{(2N+1)\sqrt{2N+3}(2N+5)\sqrt{2N+7}(2N+9)} \\ &\quad + \frac{1}{\sqrt{2N+3}(2N+5)(2N+7)(2N+9)\sqrt{2N+11}}. \end{aligned}$$

Proof: For any $M \geq N + 5$, using Lemma 3.5 and arguments similar to those presented in the proof of Theorem 3.4, we have

$$\begin{aligned} \|u - P_N^2 u\|_{H_0^2(\Lambda)}^2 &= \sum_{n=N+1}^M a_n^2 + \sum_{n=M+1}^{\infty} a_n^2 \\ &\leq \left(\frac{|\Lambda|}{2}\right)^4 \sigma_{N,M} \sum_{n=N-3}^M b_n^2 + \tilde{C}(|\Lambda|, M)^2 \sum_{n=M-3}^{\infty} b_n^2 \\ &\leq \left(\left(\frac{|\Lambda|}{2}\right)^4 \sigma_{N,M} + \tilde{C}(|\Lambda|, M)^2 \right) |u|_{H^4(\Lambda)}^2. \end{aligned}$$

For arbitrary $\varepsilon > 0$, there exists an M such that $\tilde{C}(|\Lambda|, M)^2 < \varepsilon$ and $d_i(M) < c_j(N)$, ($i = 1, 2, j = 1, 2, 3$). We now fix such an M . Then, we have $\sigma_{N,M} = \max\{c_1(N), c_2(N), c_3(N)\}$. From the definition of $c_i(N)$ in Lemma 3.5, it is easily seen that

$$\begin{aligned} N = 3 &\implies c_2(N) < c_1(N), & c_3(N) < c_1(N), \\ 4 \leq N \leq 19 &\implies c_1(N) < c_2(N), & c_3(N) < c_2(N), \\ 20 \leq N &\implies c_1(N) < c_3(N), & c_2(N) < c_3(N). \end{aligned}$$

Hence, setting

$$C_0(|\Lambda|, N) \equiv \begin{cases} \left(\frac{|\Lambda|}{2}\right)^2 \sqrt{c_1(N)}, & \text{if } N = 3, \\ \left(\frac{|\Lambda|}{2}\right)^2 \sqrt{c_2(N)}, & \text{if } 4 \leq N \leq 19, \\ \left(\frac{|\Lambda|}{2}\right)^2 \sqrt{c_3(N)}, & \text{if } 20 \leq N, \end{cases}$$

we have $\|u - P_N^2 u\|_{H_0^2(\Lambda)}^2 \leq (C_0(|\Lambda|, N)^2 + \varepsilon) |u|_{H^4(\Lambda)}^2$. Since ε is an arbitrary positive number, it holds that

$$\|u - P_N^2 u\|_{H_0^2(\Lambda)} \leq C_0(|\Lambda|, N) |u|_{H^4(\Lambda)}.$$

Thus, explicit expressions of $c_j(N)$ yield the desired results. \square

Now, we also obtain the following estimates, which further improve the constant, by a computer-assisted approach.

Theorem 3.7. *For any $N \geq 3$ and $M \geq \max\{N + 5, 20\}$, there exists a constant $C_M(|\Lambda|, N) > 0$ such that*

$$\|u - P_N^2 u\|_{H_0^2(\Lambda)} \leq C_M(|\Lambda|, N) |u|_{H^4(\Lambda)}, \quad \forall u \in H_0^2(\Lambda) \cap H^4(\Lambda). \quad (24)$$

Here,

$$C_M(|\Lambda|, N) \equiv \left(\frac{|\Lambda|}{2}\right)^2 \sqrt{\max \sigma(A) + c_3(M)}, \quad (25)$$

where A is as defined in Lemma 3.5, and $c_3(M)$ is a constant given in Lemma 3.5.

Proof: By the same argument in Theorem 3.6, using slightly different estimates, we have, for arbitrary integer $L \geq M + 5$,

$$\begin{aligned}
\|u - P_N^2 u\|_{H_0^2(\Lambda)}^2 &= \sum_{n=N+1}^M a_n^2 + \sum_{n=M+1}^L a_n^2 + \sum_{n=L+1}^{\infty} a_n^2 \\
&\leq \left(\frac{|\Lambda|}{2}\right)^4 \vec{b}^T A \vec{b} + \left(\frac{|\Lambda|}{2}\right)^4 \sigma_{M,L} \sum_{n=M-3}^L b_n^2 + \tilde{C}(|\Lambda|, L)^2 \sum_{n=L-3}^{\infty} b_n^2 \\
&\leq \left(\left(\frac{|\Lambda|}{2}\right)^4 \max \sigma(A) + \left(\frac{|\Lambda|}{2}\right)^4 \sigma_{M,L} + \tilde{C}(|\Lambda|, L)^2 \right) |u|_{H^4(\Lambda)}^2.
\end{aligned}$$

Here, we used the vector \vec{b} and matrix A given in Lemma 3.5. For arbitrary $\varepsilon > 0$, there exists an integer L such that $\tilde{C}(|\Lambda|, L)^2 < \varepsilon$ and $d_i(L) < c_j(M)$, ($i = 1, 2, j = 1, 2, 3$). For such a fixed L , based on the assumption that $M \geq 20$, we have $\sigma_{M,L} = c_3(M)$. Therefore, we obtain

$$\|u - P_N^2 u\|_{H_0^2(\Lambda)}^2 \leq \left(\left(\frac{|\Lambda|}{2}\right)^4 \max \sigma(A) + \left(\frac{|\Lambda|}{2}\right)^4 c_3(M) + \varepsilon \right) |u|_{H^4(\Lambda)}^2.$$

Since ε is arbitrary, the theorem is proven. \square

Now, let $C(|\Lambda|, N)$ denote the smallest constant satisfying the estimates given by (10) in Theorem 3.4. Then, we have the following enclosure of the optimal constant.

Theorem 3.8. *Under the assumptions of Theorem 3.7, we have*

$$\left(\frac{|\Lambda|}{2}\right)^2 \sqrt{\max \sigma(A)} \leq C(|\Lambda|, N) \leq C_M(|\Lambda|, N). \quad (26)$$

Proof: In the error estimates, we take a particular $u \in H_0^2(\Lambda) \cap H^4(\Lambda)$ such that $\vec{b} \equiv (b_{N-3}, \dots, b_M)$ coincides with an eigenvector corresponding to the maximum eigenvalue of the matrix A , as well as $b_n = 0$, or other n . Then, we

have

$$\begin{aligned}
\|u - P_N^2 u\|_{H_0^2(\Lambda)}^2 &= \sum_{n=N+1}^M a_n^2 + \sum_{n=M+1}^{\infty} a_n^2 \\
&= \left(\frac{|\Lambda|}{2}\right)^4 \max \sigma(A) \sum_{n=N-3}^M b_n^2 + \sum_{n=M+1}^{\infty} a_n^2 \\
&\geq \left(\frac{|\Lambda|}{2}\right)^4 \max \sigma(A) \sum_{n=N-3}^M b_n^2 \\
&= \left(\frac{|\Lambda|}{2}\right)^4 \max \sigma(A) |u|_{H^4(\Lambda)}^2.
\end{aligned}$$

Therefore, the optimal constant $C(|\Lambda|, N)$ satisfies

$$C(|\Lambda|, N) \geq \left(\frac{|\Lambda|}{2}\right)^2 \sqrt{\max \sigma(A)},$$

which proves the theorem. \square

Numerical Verification Results

In this subsection, we present the verified intervals that enclose the optimal constant $C(|\Lambda|, N)$ computed by expression given in Theorem 3.8. We used the following environment for verified numerical computations.

Computer environment. CPU: Intel Core2 Quad Q6700, Memory: DDR2 8GB, OS: Ubuntu Linux 7.10 AMD64, Compiler: Intel Fortran 10.1, LAPACK: version 3.1.1, BLAS: Goto BLAS 1.26, Interval arithmetic: INTLIB [3].

Table 1 shows the validated computational results of the lower bound $\frac{1}{4}\sqrt{\max \sigma(A)}$ and upper bounds $C_M(|\Lambda|, N) |\Lambda|^{-2} = \frac{1}{4}\sqrt{\max \sigma(A) + c_3(M)}$ of the optimal constants $C(|\Lambda|, N) |\Lambda|^{-2}$ for $3 \leq N \leq 32$. Here, we use the parameter $M = N + 10,000$. The real numbers in each column in the table are the lower and upper bounds of intervals given in abbreviated form. For example, in case of $N = 3$, there exists an optimal constant $C(|\Lambda|, 3) |\Lambda|^{-2}$ in the interval $[0.04469616240857, 0.04469616240858]$. We also have the following L^2 and H_0^1 estimates.

Table 1: Verification results of $C(|\Lambda|, N) |\Lambda|^{-2}$.

N	$C(\Lambda , N) \Lambda ^{-2}$	N	$C(\Lambda , N) \Lambda ^{-2}$	N	$C(\Lambda , N) \Lambda ^{-2}$
3	4.46961624085 ⁸ ₇ E-02	13	1.16345321482 ⁴ ₀ E-03	23	3.8117279298 ⁹ ₀ E-04
4	1.62145975082 ⁴ ₃ E-02	14	1.00465830406 ⁴ ₀ E-03	24	3.510274719 ⁴¹ ₃₁ E-04
5	9.00958826647 ³ ₂ E-03	15	8.7692840453 ⁵ ₁ E-04	25	3.243743962 ⁹³ ₈₃ E-04
6	5.86304210222 ³ ₁ E-03	16	7.725589432 ⁵⁴ ₄₉ E-04	26	3.00689176 ⁸⁰¹ ₇₈₉ E-04
7	4.16384630084 ⁹ ₇ E-03	17	6.8611413019 ⁷ ₂ E-04	27	2.795429334 ⁵⁷ ₄₅ E-04
8	3.1289022482 ⁴⁰ ₃₈ E-03	18	6.1366503096 ⁶ ₀ E-04	28	2.605820527 ⁶⁵ ₅₂ E-04
9	2.4467169304 ⁴⁵ ₃₉ E-03	19	5.5231170046 ⁸ ₂ E-04	29	2.435127985 ⁹² ₇₉ E-04
10	1.97099111568 ⁹ ₆ E-03	20	4.99874033 ²⁰¹ ₁₉₄ E-04	30	2.280894866 ⁵¹ ₃₇ E-04
11	1.6248684798 ⁹¹ ₈₈ E-03	21	4.546862824 ⁹⁶ ₈₈ E-04	31	2.141053068 ⁶³ ₄₈ E-04
12	1.36454584426 ⁹ ₆ E-03	22	4.15457197 ⁴⁰⁶ ₃₉₇ E-04	32	2.013851350 ³⁶ ₂₀ E-04

Theorem 3.9. *Let $C_M(|\Lambda|, N) > 0$ be the constant given in Theorem 3.7. Then, it holds that*

$$\|u - P_N^2 u\|_{L^2(\Lambda)} \leq C_M(|\Lambda|, N) \|u - P_N^2 u\|_{H_0^2(\Lambda)}, \quad \forall u \in H_0^2(\Lambda), \quad (27)$$

$$\|u - P_N^2 u\|_{H_0^1(\Lambda)} \leq \sqrt{C_M(|\Lambda|, N)} \|u - P_N^2 u\|_{H_0^2(\Lambda)}, \quad \forall u \in H_0^2(\Lambda). \quad (28)$$

Proof: The estimates given in (27) are obtained by applying Aubin-Nitsche's trick. Next, for arbitrary $u \in H_0^2(\Lambda)$, using (27), observe that

$$\begin{aligned} \|u - P_N^2 u\|_{H_0^1(\Lambda)}^2 &= \left(\frac{d}{dx}(u - P_N^2 u), \frac{d}{dx}(u - P_N^2 u) \right)_{L^2(\Lambda)} \\ &= \left(\frac{d^2}{dx^2}(u - P_N^2 u), u - P_N^2 u \right)_{L^2(\Lambda)} \\ &\leq \|u - P_N^2 u\|_{H_0^2(\Lambda)} \|u - P_N^2 u\|_{L^2(\Lambda)} \\ &\leq C_M(|\Lambda|, N) \|u - P_N^2 u\|_{H_0^2(\Lambda)}^2, \end{aligned}$$

which implies (28). \square

4. Error estimates for the finite element method in the one-dimensional case

In this section, applying Theorem 3.7 and Theorem 3.9, we derive the constructive a priori error estimates for the finite element method on one-dimensional intervals. Let Ω be a finite interval $\Omega = (\omega_0, \omega_1)$, ($\omega_0 < \omega_1$) on \mathbb{R} . Let $\omega_0 = x_0 < x_1 < \dots < x_k = \omega_1$ be a mesh of Ω and set $\Omega_i = (x_{i-1}, x_i)$. In addition, we set $h_i \equiv |\Omega_i| = x_i - x_{i-1}$ and $h \equiv (h_1, \dots, h_k) \in \mathbb{R}^k$. For an integer vector $N = (N_1, \dots, N_k) \in \mathbb{Z}^k$ with ($N_i \geq 3$), let $S_{h,N}$ be a finite-dimensional subspace of $H^2(\Omega)$ constituted of piecewise polynomials of degree N_i on Ω_i . Then, $S_{h,N}$ is generated by two types of bases, namely, a piecewise cubic Hermite polynomial whose support is two consecutive elements and a function whose support is a single element corresponding to a polynomial of degree ≥ 4 that satisfies (8).

Definition 4.1 (Hermite interpolation). Let Π_h denote a cubic Hermite interpolation from $H^2(\Omega)$ to $S_{h,N}$. That is, for each $u \in H^2(\Omega)$, $\Pi_h u \in S_{h,N}$ satisfies

$$u(x_i) = \Pi_h u(x_i), \quad \frac{du}{dx}(x_i) = \frac{d\Pi_h u}{dx}(x_i), \quad \forall i = 0, \dots, k. \quad (29)$$

Definition 4.2 (H_0^2 -projection). Let $P_{h,N}^2$ denote an H_0^2 -projection from $H_0^2(\Omega)$ to $S_{h,N}$. That is, for each $u \in H_0^2(\Omega)$, $P_{h,N}^2 u \in S_{h,N}$ is defined as

$$(u - P_{h,N}^2 u, v_{h,N})_{H_0^2(\Omega)} = 0, \quad \forall v_{h,N} \in S_{h,N}. \quad (30)$$

It follows that the Definition 4.2 is well defined, because $(\cdot, \cdot)_{H_0^2(\Omega)}$ is a bounded and coercive bilinear form on $S_{h,N}$, and Definition 4.2 ensures the unique existence of $P_{h,N}^2 u$ satisfying (30). Moreover, for each $v \in H^2(\Omega)$ with $v|_{\Omega_i} \in H_0^2(\Omega_i)$, we define $P_{N_i} v \in S_{h,N}$ such that $\text{supp } P_{N_i} v = \overline{\Omega_i}$ and $P_{N_i} v|_{\Omega_i}$ is a polynomial of degree N_i on Ω_i that satisfies (9) for $\Lambda \equiv \Omega_i$.

Theorem 4.3. Let $C_M(\cdot, \cdot)$ denote the positive constant defined in Theorem 3.7. Then, we have the following a priori error estimates for the H_0^2 -projection:

$$\|u - P_{h,N}^2 u\|_{H_0^2(\Omega)} \leq \max_{1 \leq i \leq k} C_M(h_i, N_i) |u|_{H^4(\Omega)}, \quad \forall u \in H_0^2(\Omega) \cap H^4(\Omega). \quad (31)$$

Proof: For each $u \in H_0^2(\Omega) \cap H^4(\Omega)$, from (30), we have

$$\begin{aligned} \|u - P_{h,N}^2 u\|_{H_0^2(\Omega)}^2 &= (u - P_{h,N}^2 u, u)_{H_0^2(\Omega)} \\ &= \left(u - P_{h,N}^2 u, u - \Pi_h u - \sum_{i=1}^k P_{N_i}^2 (u - \Pi_h u) \right)_{H_0^2(\Omega)}. \end{aligned}$$

Therefore,

$$\|u - P_{h,N}^2 u\|_{H_0^2(\Omega)} \leq \left\| u - \Pi_h u - \sum_{i=1}^k P_{N_i}^2 (u - \Pi_h u) \right\|_{H_0^2(\Omega)}.$$

Setting $\tilde{u} := u - \Pi_h u$, note that $P_{N_i}^2 \tilde{u}$ is uniquely determined by the definition of Π_h . In addition, taking into account that the support of $P_{N_i}^2 \tilde{u}$ coincides with $\bar{\Omega}_i$, we have

$$\left\| \tilde{u} - \sum_{i=1}^k P_{N_i}^2 \tilde{u} \right\|_{H_0^2(\Omega)}^2 = \sum_{i=1}^k \left\| \tilde{u} - P_{N_i}^2 \tilde{u} \right\|_{H_0^2(\Omega_i)}^2.$$

Thus, by Theorem 3.7, we have

$$\begin{aligned} \sum_{i=1}^k \left\| \tilde{u} - P_{N_i}^2 \tilde{u} \right\|_{H_0^2(\Omega_i)}^2 &\leq \sum_{i=1}^k C_M(h_i, N_i)^2 |\tilde{u}|_{H^4(\Omega_i)}^2 \\ &= \sum_{i=1}^k C_M(h_i, N_i)^2 \left\| \left(\frac{d}{dx} \right)^4 (u - \Pi_h u) \right\|_{L^2(\Omega_i)}^2 \\ &= \sum_{i=1}^k C_M(h_i, N_i)^2 \|u''''\|_{L^2(\Omega_i)}^2 \\ &\leq \max_{1 \leq i \leq k} C_M(h_i, N_i)^2 |u|_{H^4(\Omega)}^2. \end{aligned}$$

Here, since $(\Pi_h u)'''' = 0$, we obtain the estimates (31). \square

We also have the following L^2 and H_0^1 error estimates.

Theorem 4.4. *Under the same assumption given in Theorem, the following error estimates hold:*

$$\|u - P_{h,N}^2 u\|_{L^2(\Omega)} \leq \max_{1 \leq i \leq k} C_M(h_i, N_i) \|u - P_{h,N}^2 u\|_{H_0^2(\Omega)}, \quad \forall u \in H_0^2(\Omega), \quad (32)$$

$$\|u - P_{h,N}^2 u\|_{H_0^1(\Omega)} \leq \max_{1 \leq i \leq k} \sqrt{C_M(h_i, N_i)} \|u - P_{h,N}^2 u\|_{H_0^2(\Omega)}, \quad \forall u \in H_0^2(\Omega). \quad (33)$$

Since the proof of Theorem 4.4 is similar to that of Theorem 3.9, it is not presented in the present paper.

Remark 4.5. *Theorems 4.3 and 4.4 indicate a refinement of the estimates given in [7]. Namely, as shown in Table 1, the values of $C(|\Lambda|, N) |\Lambda|^{-2}$ obtained in the present study are approximately half that of the constant in the error estimates for the cubic Hermite interpolation ($1/\pi^2 \approx 0.101321$) presented in [7]*

Remark 4.6. *For the two-dimensional case, several constructive error estimates for H_0^2 -projection on a rectangular domain are presented in [2], [6]. Since, in these studies, the error estimates are used for the one-dimensional case, which error estimation can be improved by applying Theorem 4.3.*

Acknowledgment. The authors would like to thank Professor Siegfried M. Rump for his useful suggestions on the guaranteed computation of the upper and lower bounds of maximum eigenvalues for a symmetric matrix.

References

- [1] P.G. Ciarlet and J.L. Lions, *Handbook of Numerical Analysis Volume 5: Techniques of Scientific Computing (Part 2)*, North-Holland, Amsterdam (1997).
- [2] K. Hashimoto, K. Nagatou and M.T. Nakao, *A computational approach to constructive a priori error estimate for finite element approximations of bi-harmonic problems in nonconvex polygonal domains*, Information, **9** (2006), No. 4, 573–580.

- [3] R. Baker Kearfott, *Algorithm 763: INTERVAL ARITHMETIC: A Fortran 90 Module for an Interval Data Type*, ACM Trans. Math. Software, Vol. **22** (1996), No. 4, 385–392.
- [4] S. Kimura and N. Yamamoto, *On explicit bounds in the error for the H_0^1 -projection into piecewise polynomial spaces*, Bulletin of Informatics and Cybernetics, Vol. **31** (1999), No. 2, 109–115.
- [5] M.T. Nakao, N. Yamamoto and S. Kimura, *On the Best Constant in the Error Bound for the H_0^1 -Projection into Piecewise Polynomial Spaces*, Journal of Approximation Theory, **93** (1998), 491–500.
- [6] M.T. Nakao, K. Hashimoto and K. Nagatou, *A computational approach to constructive a priori and a posteriori error estimates for finite element approximations of bi-harmonic problems*, GAKUTO International Series, Mathematical Sciences and Applications Vol. **28**, Proceedings of the 4th JSIAM-SIMAI Seminar on Industrial and Applied Mathematics, May 26–28, 2005, Hayama, Japan (2008), 139–148.
- [7] M.H. Schultz, *Spline Analysis*, Prentice-Hall, (1973).
- [8] Y. Watanabe and M.T. Nakao, *Numerical Verifications of Solutions for Nonlinear Elliptic Equations*, Japan Journal of Industrial and Applied Mathematics, **10** (1993), No. 1, 165–178.
- [9] Y. Watanabe, M. Plum and M.T. Nakao, *A computer-assisted instability proof for the Orr-Sommerfeld problem with Poiseuille flow*, Zeitschrift fuer Angewandte Mathematik und Mechanik(ZAMM) 89 (2009), 5-18.

List of MI Preprint Series, Kyushu University

The Global COE Program
Math-for-Industry Education & Research Hub

MI

- MI2008-1 Takahiro ITO, Shuichi INOKUCHI & Yoshihiro MIZOGUCHI
Abstract collision systems simulated by cellular automata
- MI2008-2 Eiji ONODERA
The initial value problem for a third-order dispersive flow into compact almost Hermitian manifolds
- MI2008-3 Hiroaki KIDO
On isosceles sets in the 4-dimensional Euclidean space
- MI2008-4 Hirofumi NOTSU
Numerical computations of cavity flow problems by a pressure stabilized characteristic-curve finite element scheme
- MI2008-5 Yoshiyasu OZEKI
Torsion points of abelian varieties with values in infinite extensions over a p -adic field
- MI2008-6 Yoshiyuki TOMIYAMA
Lifting Galois representations over arbitrary number fields
- MI2008-7 Takehiro HIROTSU & Setsuo TANIGUCHI
The random walk model revisited
- MI2008-8 Silvia GANDY, Masaaki KANNO, Hirokazu ANAI & Kazuhiro YOKOYAMA
Optimizing a particular real root of a polynomial by a special cylindrical algebraic decomposition
- MI2008-9 Kazufumi KIMOTO, Sho MATSUMOTO & Masato WAKAYAMA
Alpha-determinant cyclic modules and Jacobi polynomials

- MI2008-10 Sangyeol LEE & Hiroki MASUDA
Jarque-Bera Normality Test for the Driving Lévy Process of a Discretely Observed Univariate SDE
- MI2008-11 Hiroyuki CHIHARA & Eiji ONODERA
A third order dispersive flow for closed curves into almost Hermitian manifolds
- MI2008-12 Takehiko KINOSHITA, Kouji HASHIMOTO and Mitsuhiro T. NAKAO
On the L^2 a priori error estimates to the finite element solution of elliptic problems with singular adjoint operator
- MI2008-13 Jacques FARAUT and Masato WAKAYAMA
Hermitian symmetric spaces of tube type and multivariate Meixner-Pollaczek polynomials
- MI2008-14 Takashi NAKAMURA
Riemann zeta-values, Euler polynomials and the best constant of Sobolev inequality
- MI2008-15 Takashi NAKAMURA
Some topics related to Hurwitz-Lerch zeta functions
- MI2009-1 Yasuhide FUKUMOTO
Global time evolution of viscous vortex rings
- MI2009-2 Hidetoshi MATSUI & Sadanori KONISHI
Regularized functional regression modeling for functional response and predictors
- MI2009-3 Hidetoshi MATSUI & Sadanori KONISHI
Variable selection for functional regression model via the L_1 regularization
- MI2009-4 Shuichi KAWANO & Sadanori KONISHI
Nonlinear logistic discrimination via regularized Gaussian basis expansions
- MI2009-5 Toshiro HIRANOUCI & Yuichiro TAGUCHI
Flat modules and Groebner bases over truncated discrete valuation rings

- MI2009-6 Kenji KAJIWARA & Yasuhiro OHTA
Bilinearization and Casorati determinant solutions to non-autonomous 1+1 dimensional discrete soliton equations
- MI2009-7 Yoshiyuki KAGEI
Asymptotic behavior of solutions of the compressible Navier-Stokes equation around the plane Couette flow
- MI2009-8 Shohei TATEISHI, Hidetoshi MATSUI & Sadanori KONISHI
Nonlinear regression modeling via the lasso-type regularization
- MI2009-9 Takeshi TAKAISHI & Masato KIMURA
Phase field model for mode III crack growth in two dimensional elasticity
- MI2009-10 Shingo SAITO
Generalisation of Mack's formula for claims reserving with arbitrary exponents for the variance assumption
- MI2009-11 Kenji KAJIWARA, Masanobu KANEKO, Atsushi NOBE & Teruhisa TSUDA
Ultradiscretization of a solvable two-dimensional chaotic map associated with the Hesse cubic curve
- MI2009-12 Tetsu MASUDA
Hypergeometric q -functions of the q -Painlevé system of type $E_8^{(1)}$
- MI2009-13 Hidenao IWANE, Hitoshi YANAMI, Hirokazu ANAI & Kazuhiro YOKOYAMA
A Practical Implementation of a Symbolic-Numeric Cylindrical Algebraic Decomposition for Quantifier Elimination
- MI2009-14 Yasunori MAEKAWA
On Gaussian decay estimates of solutions to some linear elliptic equations and its applications
- MI2009-15 Yuya ISHIHARA & Yoshiyuki KAGEI
Large time behavior of the semigroup on L^p spaces associated with the linearized compressible Navier-Stokes equation in a cylindrical domain

- MI2009-16 Chikashi ARITA, Atsuo KUNIBA, Kazumitsu SAKAI & Tsuyoshi SAWABE
Spectrum in multi-species asymmetric simple exclusion process on a ring
- MI2009-17 Masato WAKAYAMA & Keitaro YAMAMOTO
Non-linear algebraic differential equations satisfied by certain family of elliptic functions
- MI2009-18 Me Me NAING & Yasuhide FUKUMOTO
Local Instability of an Elliptical Flow Subjected to a Coriolis Force
- MI2009-19 Mitsunori KAYANO & Sadanori KONISHI
Sparse functional principal component analysis via regularized basis expansions and its application
- MI2009-20 Shuichi KAWANO & Sadanori KONISHI
Semi-supervised logistic discrimination via regularized Gaussian basis expansions
- MI2009-21 Hiroshi YOSHIDA, Yoshihiro MIWA & Masanobu KANEKO
Elliptic curves and Fibonacci numbers arising from Lindenmayer system with symbolic computations
- MI2009-22 Eiji ONODERA
A remark on the global existence of a third order dispersive flow into locally Hermitian symmetric spaces
- MI2009-23 Stjepan LUGOMER & Yasuhide FUKUMOTO
Generation of ribbons, helicoids and complex scherk surface in laser-matter Interactions
- MI2009-24 Yu KAWAKAMI
Recent progress in value distribution of the hyperbolic Gauss map
- MI2009-25 Takehiko KINOSHITA & Mitsuhiro T. NAKAO
On very accurate enclosure of the optimal constant in the a priori error estimates for H_0^2 -projection